Packaged with Tamper-Evident bottle cap. Do Not Use if breakable ring is separated or missing.

Robitussin *Honeu*



ADUIT

ADULT

ADULT

Robitussin Robitussin



DEXTROMETHORPHAN HBr (Cough Suppressant)
DOXYLAMINE SUCCINATE (Antihistamine)

MAXIMUM STRENGTH

- ✓ Controls Cough
- ✓ Relieves runny nose and sneezing







For Ages 12+ 4 FL OZ (118 mL)











ADULT

Robitussin



Active ingredients Purposes (in each 20 mL)
Dextromethorohan HBr 30 mgCouch suppressant

Dextromethorphan HBr 30 mgCough suppressant Doxylamine succinate 12.5 mg Antihistamine

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose sneezing itchy, watery eyes
- itching of the nose or throat
 controls the impulse to cough to help you sleep

Warnings

Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- trouble urinating due to an enlarged prostate gland
 glaucoma
- a cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

do not use more than directed

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- do not take more than 4 doses in any 24-hour period
 this adult product is not intended for use in children
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	20 mL every 6 hours
children under 12 years	do not use

Other information

- each 20 mL contains: sodium 21 mg
- store at 20-25°C (68-77°F).

Inactive ingredients anhydrous citric acid, blueberry juice concentrate, carboxymethylcellulose sodium, glycerin, lactic acid, natural and artificial flavors, natural grade A honey, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium gluconate, sucralose, xanthan gum, zinc gluconate

Questions or comments? call weekdays from 8 AM to 6 PM EST at 1-800-245-1040

Dist. by: Haleon, Warren, NJ 07059 ©2024 Haleon or licensor.

Trademarks owned or licensed by Haleon. For most recent product information, visit www.robitussin.com

Pat. Info www.productpats.com Made in Canada

62000000211557



