

Children's

Robitussin

AGES 6 & OVER

Drug Facts

Active ingredients (in each 10 mL)

Chlorpheniramine maleate, USP 2 mg Antihistamine
Dextromethorphan HBr, USP 15 mg Cough suppressant

Purposes

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat

Warnings

Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- trouble urinating due to an enlarged prostate gland
- glaucoma
- a cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product

- do not use more than directed
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Drug Facts (continued)

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- measure only with dosing cup provided
- keep dosing cup with product
- do not take more than 4 doses in any 24-hour period

age	dose
children under 6 years	do not use
children 6 to under 12 years	10 mL every 6 hours
adults and children 12 years and over	20 mL every 6 hours

Other information

- each 10 mL contains: sodium 6 mg
- store at 20-25°C (68-77°F)

Inactive ingredients anhydrous citric acid, FD&C red no. 40, glycerin, lactic acid, natural and artificial flavors, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments? call weekdays from 8 AM to 6 PM EST at 1-800-245-1040



Dist. by: Haleon, Warren, NJ 07059
 For most recent product information, visit www.robitussin.com
 Pat. Info www.productpats.com
 Trademarks owned or licensed by Haleon.
 ©2023 Haleon or licensor.
 Made in Canada

6200000205553



3 0031 8692 13 4

Children's

Robitussin

AGES 6 & OVER

Nighttime Cough



Long-Acting

Children's

Robitussin

AGES 6 & OVER

Nighttime Cough



Long-Acting

CHLORPHENIRAMINE MALEATE (Antihistamine)
DEXTROMETHORPHAN HBr (Cough Suppressant)

Relieves:

- Cough Up to 8 Hours
- Runny Nose

Alcohol-Free

fruit punch
flavor



4 FL OZ
(118 mL)

Do not use if plastic wrap imprinted with "SEALED FOR YOUR PROTECTION" is broken or missing.

Children's Robitussin liquid is specially formulated to provide soothing action, control your child's cough plus relieve other cold symptoms.



PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org