

73D51SBC2



Robitussin

MAXIMUM STRENGTH

12 Hour Cough & Mucus Relief

EXTENDED-RELEASE TABLETS

GUAIFENESIN & DEXTROMETHORPHAN HYDROBROMIDE
1200 mg/60 mg EXTENDED-RELEASE TABLETS
Expectorant & Cough Suppressant

- ✓ **Controls Cough**
- ✓ **Thins & Loosens Mucus**



Actual Size

8

EXTENDED-RELEASE TABLETS



73D51SBC2

MAXIMUM STRENGTH
12 Hour
Cough & Mucus Relief
EXTENDED-RELEASE TABLETS

Robitussin

Do not use if blister
unit is broken or torn.**12 Hour** Cough & Mucus Relief**MAXIMUM STRENGTH****EXTENDED-RELEASE**
TABLETS

Robitussin

Drug Facts

Active ingredients **Purposes**
*(in each extended-release tablet)*Dextromethorphan HBr 60 mg Cough suppressant
Guaifenesin 1200 mg.....Expectorant**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help you get to sleep

Warnings**Do not use**

- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

When using this product

- do not use more than directed

Stop use and ask a doctor if

- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

Drug Facts (continued)**If pregnant or breast-feeding**, ask a health professional before use.**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).**Directions**

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: 1 tablet every 12 hours; not more than 2 tablets in 24 hours
- children under 12 years of age: do not use

Other information

- each tablet contains: magnesium 25 mg
- store between 20-25°C (68-77°F)

Inactive ingredients

carbomer homopolymer type B, copovidone, D&C yellow #10 aluminum lake, hypromellose, magnesium hydroxide, magnesium stearate, microcrystalline cellulose, silicon dioxide

Questions or comments? call weekdays from 8 AM to 6 PM EST at **1-800-245-1040****PARENTS:**

Learn about teen medicine abuse

www.StopMedicineAbuse.org

Keep the carton. It contains important information.
Distributed by: Haleon, Warren, NJ 07059
For most recent product information, visit www.robitussin.com
Trademarks owned or licensed by Haleon.
©2023 Haleon or licensor.
205420

3

**81% For Position Only**

0031 8765 08 1

73D51 SB C2