

NEW! ADULT

HALEON

# Robitussin

Nighttime Cough DM

Soft CHEWS

DEXTROMETHORPHAN HBr 15 mg  
(Cough Suppressant)  
DOXYLAMINE SUCCINATE 6.25 mg  
(Antihistamine)

**FAST relief ANYWHERE**

Cough, runny  
nose & sneezing

20 CHEWABLE TABLETS

Chew tablets completely before swallowing

UP  
TO 8HR  
COUGH RELIEF



Berry  
flavor

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Distributed by: Haleon, Warren, NJ 07059  
Trademarks are owned by or licensed to the Haleon group of companies.  
Pat. Info www.productipats.com ©2023 Haleon group of companies or its licensor.  
For most recent product information, visit www.robitussin.com

**PARENTS:**

Learn about teen medicine abuse

www.StopMedicineAbuse.org

208797

51141421



Peel Here

**Drug Facts**

**Active ingredients (in each chewable tablet) Purposes**

Dextromethorphan equivalent to dextromethorphan HBr 15 mg..... Cough suppressant  
Doxylamine succinate 6.25 mg..... Antihistamine

**Uses**

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - runny nose
  - sneezing
  - itchy, watery eyes
  - itching of the nose or throat
- controls the impulse to cough to help you sleep

**Drug Facts (continued)**

**Warnings**

- Do not use**
- to sedate a child or to make a child sleepy
  - if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- trouble urinating due to an enlarged prostate gland
- glaucoma
- a cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema

**Drug Facts (continued)**

**Ask a doctor or pharmacist before use if you are** taking sedatives or tranquilizers

- When using this product**
- marked drowsiness may occur
  - avoid alcoholic drinks
  - alcohol, sedatives, and tranquilizers may increase drowsiness
  - be careful when driving a motor vehicle or operating machinery
  - excitability may occur, especially in children

**Stop use and ask a doctor if** cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache.  
These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than 4 doses (8 chewable tablets) in any 24-hour period
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	take 2 chewable tablets every 6 hours, as needed
children under 12 years	do not use

**Drug Facts (continued)**

**Other information**

- each chewable tablet contains: potassium 10 mg, sodium 7 mg
- store at a controlled room temperature 20-25°C (68-77°F)

**Inactive ingredients**

carboxymethylcellulose calcium, carnauba wax, corn syrup solids, crospovidone, D&C red no. 30 aluminum lake, FD&C blue no. 1 aluminum lake, glycerin, mannitol, natural and artificial flavors, pregelatinized starch, simethicone, sodium gluconate, sorbitol, sorbitol solution, sucralose, sucrose, whole dry milk

**Questions or comments?**

call weekdays from 8 AM to 6 PM EST at 1-800-245-1040